



**O‘ZBEKISTON RESPUBLIKASI
VETERINARIYA VA CHORVACHILIKNI RIVOJLANTIRISH
QO‘MITASI
VETERINARIYA ILMIY-TADQIQOT INSTITUTI**



**“VETERINARIYA SOHASIDAGI DOLZARB
MUAMMOLAR YECHIMI YOSH
TADQIQOTCHILAR TALQINIDA”**

RESPUBLIKA ILMIY-AMALIY KONFERENSIYASI

TO‘PLAMI – 1/24

SAMARQAND – 2024

“Veterinariya sohasidagi dolzarb muammolar yechimi yosh tadqiqotchilar talqinida” Respublika ilmiy-amaliy konferensiyasi materiallari To‘plami – 1/24. Samarqand 2024-yil. 297 bet.

Ushbu to‘plamda veterinariya va qishloq xo‘jaligi sohalaridagi dolzarb muammolarini hal etishda tadqiqotchilar tomonidan olib borilayotgan ilmiy izlanishlarning ahamiyati, hayvonlarda uchrovchi turli xil yuqumli, yuqumsiz va parazitlar kasalliklarning etiologiyasi, patogenezi, diagnostikasi, davolash, oldini olish va qarshi kurash shuningdek chorva hayvonlarini saqlash hamda oziqlantirish borasida yangi va innovatsion chora-tadbirlar haqida hamda so‘ngi yillarda Veterinariya ilmiy-tadqiqot institutida yaratilgan va amaliyotga joriy qilingan diagnostikumlar, allergenlar, vaksinalar va boshqa erishilgan yutuqlar bo‘yicha ma’lumotlar keltirilgan.

To‘plam materiallaridan veterinariya va chorvachilik sohasidagi OTM va ITM larda ilmiy faoliyat bilan shug‘ullanuvchi ilmiy xodimlar, tayanch doktorant va doktorantlar, Oliy ta’lim muassalari bakalavriat hamda magistratura bosqichlarida tahsil oladigan talabalar foydalanishi mumkin.

To‘plam Veterinariya ilmiy-tadqiqot instituti Ilmiy kengashi tomonidan chop etishga tavsiya qilingan.

Eslatma: Konferensiya materiallari to‘plamiga kiritilgan maqolalarning mazmuni uchun nashriyot va tashkiliy qo‘mita javobgar emas. Maqolalar mualliflar matni asosida chop etildi.

Veterinariya ilmiy-tadqiqot instituti 2024-yil.

O.A.Djuraev, Ph.D., Senior Researcher,
A.Kh.Khushnazarov, junior research fellow
Veterinary Scientific Research Institute

Annotation: The article highlights the results of scientific research on the pathoanatomical and histological study of experimental rabbits infected with atypical mycobacteria. It has been established that the latter are able to cause rather deep pathological processes in their body. These processes are characterized, in addition to general hemodynamic disorders, by the development of necrotic foci of various sizes in the liver.

Key words: *atypical mycobacteria, rabbits, autopsy, morphology, histoexamination, histosections, internal organs.*

Аннотация: В статье освещены результаты научных исследований по патологоанатомическому и гистологическому изучению подопытных кро-ликов, зараженных атипичными микобактериями. Установлено, что послед-ние способны вызывать довольно глубокие патологические процессы в их организме. Эти процессы характеризуются, помимо общих гемодинамичес-ких расстройств, развитием некротических очагов различных размеров в печени.

Ключевые слова: *атипичные микобактерии, кролики, патвскрытие, морфология, гистоисследование, гистосрезы, внутренние органы.*

Аннотация: Мақолада туберкулёзнинг нотипик микобактериялари юктирилган қуёнлар организмнинг патологоанатомик ва гистологик текши-ришлар натижалари ёритилган. Тажриба қуёнлари организмда нотипик микобактериялар анча чуқур патологик жараёнлар келтириб чиқариши аниқ-ланган. Мазкур жараёнлар, умумий гемодинамик бузилишлардан ташқари, жигарда турли ўлчамлардаги некроз ўчоқларининг ривожланиши билан ифодаланади.

Калит сўзлар: *нотипик микобактериялар, қуёнлар, патологоанатомик ёриш, морфология, гистотекишириш, гистокесмалар, ички аъзолар.*

Relevance of the problem. It is known that livestock breeding is of great importance in improving the agricultural production of the republic, the development of which is impossible without reliable protection of animals from various diseases. However, the increase in the production of livestock products is hampered by various infectious diseases, among which tuberculosis occupies a special place. And this problem is becoming even more acute due to the fact that in recent years the purchase and import of imported highly productive precious cattle has sharply increased. In addition to all this, imported animals are very predisposed and susceptible to atypical mycobacteria, and also react positively in allergy tests for tuberculosis. The disease not only causes significant economic damage, but also has great socio-epidemiological significance.

Increasingly, reports are appearing in the scientific literature indicating the identification of a significant number of animals with “nonspecific reactions” during allergy studies. These reactions have been observed in animals infected with so-called atypical mycobacteria. Currently, more than 300 species of atypical mycobacteria are known, the isolation of which has dramatically complicated the problem of eliminating tuberculosis. This is explained by the unreasonable slaughter of highly productive animals and birds that respond positively to the introduction of tuberculin, as well as the imposition of restrictive measures on the farm, household, etc. The import of a large number of cattle to Uzbekistan from neighboring countries and Europe, where atypical mycobacteria are widespread, has further aggravated the task of veterinary specialists. Increasingly, reports are appearing in the scientific literature indicating the identification of a significant number of animals with “nonspecific reactions” during allergy studies. These reactions have been observed in animals infected with so-called atypical mycobacteria. Currently, more than 300 species of atypical mycobacteria are known, the isolation of which has dramatically complicated the problem of eliminating tuberculosis. This is explained by the unreasonable slaughter of highly productive animals and birds that respond positively to the introduction of tuberculin, as well as the imposition of restrictive measures on the

farm, household, etc. The import of a large number of cattle to Uzbekistan from neighboring countries and Europe, where atypical mycobacteria are widespread, has further aggravated the task of veterinary specialists.

Material and research methods. In order to study the pathogenic effect on the morphology of internal organs, an experimental intranasal infection was carried out with reference strains of atypical mycobacteria of rabbits, divided into 3 groups (9 animals each). The first and second groups of rabbits were infected, respectively, with a bacterial mass of *M.kansasii* and *M.intracellulare* strains at a dose of 5 mg/kg; the third group served as a control and was not infected. In order to study the dynamics of the development of pathological processes, rabbits were subjected to planned slaughter 2, 4 and 6 months after infection, three animals from each group.

At each slaughter, the pathological anatomical picture of the autopsy of all experimental animals was carefully studied. The pathological material obtained from each animal was subjected to appropriate histological processing; after staining with hematoxylin-eosin, all histological sections were examined under a microscope.

Research results. During control slaughter, to study the dynamics of the development of pathological processes in the body of experimental animals, the following changes were established:

1 – group. All rabbits killed 2 months after infection showed congestion of the lungs with small areas of atelectasis and focal pinpoint hemorrhages, as well as multiple necrotic lesions the size of millet grains on the liver with curdled contents.

4 months after the start of the experiments, the liver, spleen and lymph nodes of all rabbits were enlarged and hyperemic, and in two animals there were dense foci of gray necrosis on the lungs. In one rabbit, the entire intestinal tract, especially its thick section, is diffusely affected by necrotic foci of various sizes. The autopsy picture 6 months after infection was characterized by the discovery of numerous foci of necrosis penetrating deep into the liver parenchyma, as well as severe enlargement of the lymph nodes.

2 – group. In animals killed 2 months after infection, the liver is enlarged, on its surface in one rabbit there are lesions with small punctate sizes, and in others there are extensive areas of necrosis with curdled contents, the kidneys and spleen are swollen and stagnant. At slaughter 4 and 6 months after infection, no significant changes were detected in the internal organs.

At all times of slaughter, no pathological changes were found in rabbits of the 3-rd control group at autopsy.

Pathological material taken from all experimental laboratory animals, after appropriate histological processing, cutting and staining, was subjected to a thorough microscopic examination, the results of which are given below.

Thus, morphological changes in rabbits of group 1, killed 2 months after infection, were characterized by hemodynamic disorders in the form of hyperemia, stasis and hemorrhages in the lungs; the presence of small and extensive hemorrhages, granular degeneration, micronecrosis of hepatocytes in the liver and convoluted tubule epithelium in the kidneys; infiltration of lymphoid elements of the connective tissue stroma.

In experimental animals killed 4 months later, serous edema with diapedetic hemorrhages, infiltration of serous-hemorrhagic exudate, desquamation and necrobiosis of alveocytes and bronchial epithelium are noted in the lungs. Venous congestion, depletion of the pulp in lymphocytes, increased hemosiderosis and hemorrhages were found in the spleen. The villi of the small intestine are shortened and deformed; there is no epithelial cover on them. The sinuses of the lymph nodes are dilated and filled with exudate containing reticular cells, lymphoblasts, lymphocytes, single eosinophils and erythrocytes.

In rabbits killed 6 months after the start of the experiments, the liver had multiple foci of necrobiosis and necrosis of hepatocytes, hyalinosis of the walls of the central veins and arteries. Granular dystrophy was established in the kidneys, hyperplasia was developed in the spleen due to the growth of the reticular base, and the number of plasma cells was increased.

Microscopic studies of histosections from the internal organs of rabbits of group 2, killed 2

months after infection, were distinguished by numerous small foci of necrosis in the liver. Many of them focus on the breakdown of cell nuclei into clumps or their transformation into a homogeneous mass. Necrotic lesions are surrounded by a belt consisting of epithelioid and plasma cells. In the kidneys - dilation and congestion of blood vessels, granular degeneration and necrobiosis of the convoluted tubule epithelium.

Histological studies of the internal organs of experimental animals killed 4 and 6 months after infection did not reveal significant changes. Microscopic studies of histosections from the internal organs of rabbits of group 2, killed 2 months after infection, were distinguished by numerous small foci of necrosis in the liver. Many of them focus on the breakdown of cell nuclei into clumps or their transformation into a homogeneous mass. Necrotic lesions are surrounded by a belt consisting of epithelioid and plasma cells. In the kidneys - dilation and congestion of blood vessels, granular degeneration and necrobiosis of the convoluted tubule epithelium.

Histological studies of the internal organs of experimental animals killed 4 and 6 months after infection did not reveal significant changes.

Conclusion. Thus, the results of pathological and histological studies of experimental rabbits infected with atypical mycobacteria showed that the latter are capable of causing quite profound pathological processes in their body. These processes are characterized, in addition to general hemodynamic disorders, by the development of necrotic foci of various sizes in the liver. At the same time, the dynamics and degree of development of these changes in the structure of the organs of experimental animals are very diverse, but they are more pronounced in experimental rabbits of the first group infected with a strain of atypical mycobacteria *M.kansasii*.

LIST OF USED LITERATURE:

1. Мамадуллаев, Г., Элмуродов, Б., Джураев, О., Джуракулов, О., & Файзиев, У. (2022). Рифизостреп–новый комбинированный препарат против микобактерий туберкулёза. *in Library*, 22(2).
2. Джураев, О., & Хушназаров, А. (2023). Порядок и методы патологического обследования сельскохозяйственных животных. *in Library*, 3(3), 21-25.
3. Джураев, О. (2024). Общее определение патоморфологических изменений при инфекционных болезнях. *in Library*, 1(1), 28-31.
4. Джураев, О. (2024). Порядок анализа и обобщения результатов патологоанатомического исследования. *in Library*, 1(1), 25-26.
5. Джураев, О. (2023). Патология сельскохозяйственных животных проверка на краску. *in Library*, 4(4), 28-29.
6. Джураев, О. А. (2023). РЕЗУЛЬТАТЫ ПАТОМОРФОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ ПО ИСПЫТАНИЮ НОВОГО КОМБИНИРОВАННОГО ПРЕПАРАТА “РИФИЗОСТРЕП”. *Gospodarka i Innowacje.*, 35, 517-520.
7. Джураев, О. А., & Мамадуллаев, Г. Х. (2023, May). Влияние рифизострепа на морфологию внутренних органов. In *International Conference on Research Identity, Value and Ethics* (pp. 387-391).
8. Davlatov, R. B., & Khushnazarov, A. K. (2024). Diagnosis and chemoprophylaxis of rabbit eumeriosis. In *E3S Web of Conferences* (Vol. 480, p. 03020). EDP Sciences.
9. Газнакулов, Т. К., Орипов, А. О., Сафаров, А. А., Хушназаров, А. Х., Давлатов, Р. Б., Абдухакимов, Ш., & Мавланов, С. (2023). ХС Салимов, МК Бутаев, ЗЭ Рузиев,– Биохавфсизлик.
10. Джураев, О., & Хушназаров, А. (2023). Порядок и методы патологического обследования сельскохозяйственных животных. *in Library*, 3(3), 21-25.
11. Хушназаров, А., Абдиев, Ф., & Акрамов, К. (2019). Распространение тениаринхоза на побережье Амударьи. *in Library*, 19(3), 25-26.
12. Хушназаров, А. Х., & Акрамов, К. Ш. ТЕНИАРИНХОЗНИНГ АМУДАРЁ СОҲИЛЛАРНДА ТАРҚАЛИШИ.
13. Газнакулов, Т., & Хушназаров, А. (2023). Литературный обзор по истории развития эпизоотологии и изучения бешенства. *in Library*, 1(2), 7-9.

VETERINARIYA ILMIY-TADQIQOT INSTITUTI

“VETERINARIYA SOHASIDAGI DOLZARB MUAMMOLAR YECHIMI TADQIQOTCHILAR TALQINIDA”

RESPUBLIKA ILMIY-AMALIY KONFERENSIYASI

TO‘PLAMI – 1/24

Bosishga ruxsat etildi __.__.2024
Bichimi 210x297, «Times New Roman»
garniturası. Raqamli bosma usulida bosildi.
Adadi 100 dona. Buyurtma: № 12/24



Nashriyot guvohnomasi № 035928

“SHUHRAT MENGLIYEV” MCHJ bosmaxonasida chop etildi.
Manzil: Samarqand sh., M.Ulug‘bek ko‘chasi, 101-uy(94) 572-04-07