

PREDICTION OF DENTAL MORBIDITY IN CHEMICAL INDUSTRY WORKERS

Ibragimova F.I.

Bukhara state medical institute

Relevance. Chemicals of industrial air environment are found in oral fluid, hard tissues of teeth, dental deposits, biopsies of tissue structures. They aggravate the course of dental caries due to the substitution of calcium ions in hydroxyapatite crystals, chronic inflammation in periodontal tissues, violation of the integrity of the epithelium of the oral mucosa.

However, the influence of chemical factors of production on the oral cavity of workers, the consequences of these effects have not been fully studied. Given this, the need to continue dental, clinical, functional and microbiological research on this problem has not lost its relevance.

The purpose of the study The aim is to develop prevention and improve the prediction of dental diseases based on the proposed conceptual approach in chemical industry workers.

The object of the study. The study will examine 4,150 chemical industry workers aged 25 to 60 years who have had direct contact with chemical elements for several years. All the surveyed will be divided into 3 groups: 1 - group - workshop workers, these are persons directly exposed to chemical elements; 2-group - locksmiths, plumbers, installers, cleaners, laboratory assistants, persons who rarely come into contact with chemical compounds; 3 - group – industrial administration, who have no contact with chemical elements.

Dental examination of workers with harmful chemical working conditions showed that the intensity of the carious process, the prevalence of periodontal diseases, pathological erasability, dentin hypersensitivity, the predominance of precancerous diseases among the diseases of the SOPR, the low level of oral hygiene is higher compared to the residents of the city. The high level of dental morbidity is due to the presence of harmful factors of chemical production, the lack of planned sanitation of the oral cavity, an effective system of prevention and medical examination at industrial enterprises.

At all three chemical production enterprises, with increasing work experience, the intensity of the carious process decreases, the number of teeth removed increases; the number of people with precancerous diseases of the SOPR and periodontal diseases increases. SOPR diseases at the early stages of diagnosis do not always have pronounced clinical symptoms, but are characterized by obvious pathological changes in the epithelium, confirmed morphologically.

A step-by-step method of primary and secondary prevention of dental diseases in chemical industry workers has been developed and recommended for dental practice. The indicators of the need for dental treatment among workers of harmful production factors are determined, and the volume of necessary complex treatment among workers of the chemical industry is calculated. The results obtained are planned to be included in the medical practice of institutions

conducting experimental research, as a methodological recommendation, a patent for an invention, scientific and innovative work.

Dental morbidity of workers of chemical enterprises, identification of harmful environmental factors and chemical production, reorganization of the dental care system, introduction of dental hygienist positions in the staffing of the dental service of medical units of chemical plants, timely dispensary observation, annual screening of the condition of the mucous membrane of the cavity for oncopathology, prevention of dental diseases and active sanitary and educational work and systematic promotion of a healthy lifestyle. Identify the main risk factors and form risk groups among the population in need of dynamic monitoring of dental health and systematic prevention and medical examination. Carrying out planned prevention of dental diseases, increasing the level of individual oral hygiene of workers, it is necessary to reorganize dental services of harmful industries and include dental hygienists in their structure.

In order to reduce the prevalence and intensity of major dental diseases and improve the hygienic condition of the oral cavity in workers of chemical enterprises, it is necessary to carry out preventive measures, including dental education, brushing teeth with multicomponent therapeutic and preventive toothpastes, promotion of a healthy lifestyle.

References

1. АРИФОВ, СС, and МБ ТУХТАЕВ. "Учредители: Общество с ограниченной ответственностью с участием иностранного капитала" E-LINE PRESS." *СТОМАТОЛОГИЯ Учредители: Общество с ограниченной ответственностью с участием иностранного капитала* E-LINE PRESS" 1: 82-85.

2. Юнусходжаева, М., and Л. Хасанова. "НАРУШЕНИЕ АГРЕГАЦИОННОЙ ФУНКЦИИ ТРОМБОЦИТОВ У ПАЦИЕНТОВ С БЫСТРОПРОГРЕССИРУЮЩИМ ПАРОДОНТИТОМ В РАЗНЫХ СТЕПЕНЯХ ТЯЖЕСТИ." *Медицина и инновации* 1.3 (2021): 99-102.

3. Даминова, Ш., et al. "Study of the influence of obesity in parentals on teething in children of the first year of life who are exclusively breastfed." in *Library* 22.1 (2022): 86-88.

4. Воробьев, Анатолий Андреевич, et al. "Медицинская микробиология, вирусология и иммунология." (2015).